

Sex differences in brain activation during a working memory task in adult patients with ADHD

Poster Number:

2486

Submission Type:

Abstract Submission

Authors:

Peter Soros¹, Katharina Bachmann¹, Alexandra Lam^{1,2}, Manuela Kanat³, Eliza Hoxhaj⁴, Swantje Matthies⁴, Bernd Feige⁴, Alexandra Philipsen^{1,4}

Institutions:

¹Psychiatry and Psychotherapy - University Hospital, University of Oldenburg, Oldenburg, Germany, ²Department of Psychology, Biological Psychology Lab, Medical Campus University of Oldenburg, School of Medicine and Health Sciences, University of Oldenburg, Oldenburg, Germany, ³Department of Psychology, Laboratory for Biological Psychology, University of Freiburg, Freiburg, Germany, ⁴Department of Psychiatry and Psychotherapy, University of Freiburg, Freiburg, Germany

Introduction:

Attention-deficit/hyperactivity disorder (ADHD) in adulthood is a serious and frequent psychiatric disorder with a strong neurobiological background (Bachmann et al. 2016). Task-based fMRI, using a variety of sensory and cognitive paradigms, has demonstrated hypoactivation of a fronto-parietal network and hyperactivation of visual, dorsal attention and default mode networks (Cortese et al. 2012). Most participants in these neuroimaging studies were men and thus it remains unclear if changes in brain activation found in men with ADHD also apply to women with ADHD. To study potential sex differences in neural function, we acquired fMRI during a one-back working memory task in women and men with ADHD.

Methods:

Brain imaging data from 31 women (42±11 years) and 24 men (37±12 years) with adult ADHD were analyzed. The diagnosis of ADHD was established by expert psychiatrists and validated using observer-rating and self-rating scales. The study was approved by the local ethics committee.

To probe working memory function, patients performed a one-back letter task during fMRI. With button presses of their left and right hand, patients had to indicate if the presented letter was a target (preceding letter was the same) or non-target (preceding letter was different). Patients performed 2 runs of 300 stimuli (89 target, 211 non-target; interstimulus interval: 1 s) each.


T1-weighted anatomical and T2*-weighted functional images were acquired on a Siemens Magnetom Trio at 3 T. For BOLD fMRI, gradient-echo echoplanar imaging was performed (TR: 2250 ms, TE: 30 ms, voxel size: 3x3x3 mm³, 36 slices).

fMRI data analysis was carried out using FSL. Pre-processing included motion correction, non-brain removal, spatial smoothing (5 mm) and highpass temporal filtering. A linear model was then fitted to the fMRI data. Six events were modeled as regressors of interest: correct and incorrect responses as well as omissions for target and non-target stimuli. Individual age was added as regressor of no interest in the group analysis.

Results:

Behavioral performance during the fMRI experiment was not significantly different between women and men. Women identified 61±14% of target trials correctly, men 60±16% (p=0.83). Reaction time in target-correct trials was 573±73 ms in women and 546±73 ms in men (p=0.2).

Brain activation associated with target-correct trials was significantly stronger in men compared to women in following areas (Figure): bilateral posterior cingulate gyrus, left superior parietal lobule, right frontal pole and right lentiform nucleus. Brain areas with significantly stronger activation in women compared to men were not found.



Brain activation map, men > women. Brain regions color-coded in red and yellow show stronger activation in men than in women. Z statistic images are overlaid onto the mean structural T1-weighted image of the participants. The upper two rows show axial images of the brain in radiological convention (the right hemisphere is on the left). The last row shows sagittal images of the right (images 1-3) and left hemisphere (image 4). LN: lentiform nucleus. P: putamen. FP: frontal pole. PCG: posterior cingulate gyrus. SPL: superior parietal lobule.

Conclusions:

We performed a direct comparison of brain activation during a working memory task between men and women with ADHD. While the accuracy of responses and the latency of correct responses was not significantly different between the sexes, men showed several clusters of increased neural activation relative to women. The largest cluster contained the bilateral posterior cingulate gyrus, a key component of the default mode network (Greicius et al. 2013). The default mode network is a set of brain areas that is active during rest and that deactivates during a task. Activation of the posterior cingulate gyrus during the one-back task may suggest a dysfunction of the default mode network in men with ADHD. In addition, men had greater brain activation in frontal and striatal regions that, while subserving many different functions, are an important part of the circuits generating executive functions. Our results imply that men recruit larger neural resources than women to achieve similar behavioral performance during the one-back task. In summary, our study adds to the small corpus of literature on sex differences in adult ADHD (Valera et al. 2010) and provides evidence that cognitive functions may be represented differently in men and women with ADHD.

Disorders of the Nervous System:

Other Psychiatric Disorders ¹

Imaging Methods:

BOLD fMRI ²

Learning and Memory:

Working Memory

Keywords:

Attention Deficit Disorder
FUNCTIONAL MRI
Memory
Psychiatric
Sexual Dimorphism
Other - ADHD

¹²Indicates the priority used for review

Would you accept an oral presentation if your abstract is selected for an oral session?

Yes

I would be willing to discuss my abstract with members of the press should my abstract be marked newsworthy:

Yes

Please indicate below if your study was a "resting state" or "task-activation" study.

Task-activation

By submitting your proposal, you grant permission for the Organization for Human Brain Mapping (OHBM) to distribute the presentation in any format, including video, audio print and electronic text through OHBM OnDemand, social media channels or other electronic media and on the OHBM website.

I accept

Healthy subjects only or patients (note that patient studies may also involve healthy subjects):

Patients

Internal Review Board (IRB) or Animal Use and Care Committee (AUCC) Approval. Please indicate approval below. Please note: Failure to have IRB or AUCC approval, if applicable will lead to automatic rejection of abstract.

Yes, I have IRB or AUCC approval

Please indicate which methods were used in your research:

Functional MRI

For human MRI, what field strength scanner do you use?

3.0T

Which processing packages did you use for your study?

FSL

Provide references in author date format

Bachmann, K. et al. (2016) 'Mindfulness-Based cognitive therapy and the Adult ADHD Brain: A Neuropsychotherapeutic Perspective', *Frontiers in Psychiatry*, vol. 7, article 117.

Cortese, S. et al. (2012), 'Toward Systems Neuroscience of ADHD: A Meta-analysis of 55 fMRI Studies', *American Journal of Psychiatry*, vol. 169, pp. 1038-1055.

Greicius, M.D. et al. (2013), 'Functional Connectivity in the Resting Brain: A Network Analysis of the Default Mode Hypothesis', *Proceedings of the National Academy of Sciences*, vol. 100, pp. 253-258.

Valera, E.M. et al. (2010), 'Sex Differences in the Functional Neuroanatomy of Working Memory in Adults With ADHD', *American Journal of Psychiatry*, vol. 167, pp. 86-94.